

- Ozcan, H.K., Ucan, O.N., Sahin, U., Borat, M. and Bayat, C. (2006), "Artificial neural network modeling of methane emissions at Istanbul Kemerburgaz-Odayeri landfill site", *J. Sci. Ind. Res.*, **65**(2), 128 p.
- Ruiz-Suarez, J.C., Mayora-Ibarra, O.A., Torres-Jimenez, J. and Ruiz-Suarez, L.G. (1995), "Short-term ozone forecasting by artificial neural networks", *Adv. Eng. Software*, **23**(3), 143-149.
- Tasadduq, I., Rehman, S. and Bubshait, K. (2002), "Application of neural networks for the prediction of hourly mean surface temperatures in Saudi Arabia", *Renew. Energy*, **25**(4), 545-554.
- US.EPA (1993), Integrated Risk Information System on Ethylbenzene, Cincinnati, OH, USA, P 36, 99, 124.
- Viotti, P., Liuti, G. and Genova, P.Di. (2002), "Atmospheric urban pollution: Applications of an artificial neural network (ANN) to the city of Perugia", *Ecol. Model.*, **148**(1), 27-46.
- Voukantsis, D., Karatzas, K., Kukkonen, J., Räsänen, T., Karppinen, T. and Kolehmainen, M. (2011), "Inter comparison of air quality data using principal component analysis, and forecasting of PM₁₀ and PM_{2.5} concentrations using artificial neural networks, in Thessaloniki and Helsinki", *Sci. Total Environ.*, **409**(7), 1266-1276.
- Wu, W., Dandy, G.C. and Maier, H.R. (2014), "Protocol for developing ANN models and its application to the assessment of the quality of the ANN model development process in drinking water quality modelling", *Environ. Model. Software J.*, **54**, 108-127.
- Yi, J. and Prybutok, V.R. (1996), "A neural network model forecasting for prediction of daily maximum ozone concentration in an industrialized urban area", *Environ. Pollut.*, **92**(3), 349-357.

WL